Pre/post operation

Cutting through all the information

Due to illness, disease or trauma, your pet may one day require surgery. While potentially stressful (for both you and your pet) there are a few basic guidelines that you can follow that will make the process as complication-free as possible and put your pet on the fast road to recovery.

Depending on the type of surgery, whether minor or major, your veterinarian will advise you when your pet can resume his/her normal lifestyle.

Pre-surgical instructions

- Your veterinarian will do a check-up on your pet before the surgery to determine if there are any pre-existing conditions that may interfere with the surgical procedure.
- Make sure your pet is current on his/her vaccinations.
- Your veterinarian may suggest a blood test to screen for disease not apparent from a physical examination.
- No food after
- No water after



Post-surgical instructions

- Chances are your pet will be weak or groggy after surgery. Do not let him/her get too excited.
- Restrain your animal with a leash or put them in a carrier when leaving the clinic. This will protect them from additional injury.
- Provide only small amounts of food and water until he/she readjusts to • being at home and is recovering. Too much food and water can lead to upset stomachs or vomiting.
- If a special post-surgical diet has been prescribed, follow all instructions carefully.
- Limit your pet's exercise. Climbing stairs, jumping or running may open up sutures or cause nausea.
- Make sure his/her sleeping area is clean, warm and free of draughts.
- Your veterinarian may prescribe medication to administer during your pet's recovery. Follow all label instructions carefully.
- Sutures are usually removed approximately 10 days after surgery. Check the area around the incision daily for any redness, swelling or discharge. If you detect any irritation, contact your veterinarian immediately.
- Prevent your pet from licking or chewing on the wound. If this is difficult to do, you should provide a physical barrier by placing an 'Elizabethan collar' around their head.

/

Suture removal due on 1



